

Research article

Open Access

## Quantitative trait loci in *Anopheles gambiae* controlling the encapsulation response against *Plasmodium cynomolgi* Ceylon

Liangbiao Zheng\*<sup>1</sup>, Shuang Wang<sup>1</sup>, Patricia Romans<sup>2</sup>, Hongyu Zhao<sup>1</sup>, Coralia Luna<sup>1</sup> and Mark Q Benedict<sup>3</sup>

Address: <sup>1</sup>Yale University School of Medicine, Department of Epidemiology and Public Health, 60 College Street, New Haven, CT 06520, USA, <sup>2</sup>Department of Zoology, University of Toronto, Toronto, ON, Canada M5S 3G5 and <sup>3</sup>Centers for Disease Control and Prevention, Mailstop F22, Chamblee, GA 30334, USA

Email: Liangbiao Zheng\* - liangbiao.zheng@yale.edu; Shuang Wang - shuang.wang@yale.edu; Patricia Romans - victorie@zoo.utoronto.ca; Hongyu Zhao - hongyu.zhao@yale.edu; Coralia Luna - coralia.luna@yale.edu; Mark Q Benedict - mqb0@cdc.gov

\* Corresponding author

Published: 24 October 2003

Received: 10 July 2003

BMC Genetics 2003, 4:16

Accepted: 24 October 2003

This article is available from: <http://www.biomedcentral.com/1471-2156/4/16>

© 2003 Zheng et al; licensee BioMed Central Ltd. This is an Open Access article: verbatim copying and redistribution of this article are permitted in all media for any purpose, provided this notice is preserved along with the article's original URL.

### Abstract

**Background:** *Anopheles gambiae* females are the world's most successful vectors of human malaria. However, a fraction of these mosquitoes is refractory to *Plasmodium* development. L3-5, a laboratory selected refractory strain, encapsulates transforming ookinetes/early oocysts of a wide variety of *Plasmodium* species. Previous studies on these mosquitoes showed that one major (*Pen1*) and two minor (*Pen2*, *Pen3*) autosomal dominant quantitative trait loci (QTLs) control the melanotic encapsulation response against *P. cynomolgi* B, a simian malaria originating in Malaysia.

**Results:** We have investigated the response of L3-5 to infection with *P. cynomolgi* Ceylon, a different but related parasite species, in crosses with the susceptible strain 4Arr. Refractoriness to this parasite is incompletely recessive. Infection and genotyping of F2 intercross females at genome-spanning microsatellite loci revealed that 3 autosomal QTLs control encapsulation of this species. Two loci map to the regions containing *Pen2* and *Pen3*. The novel QTL maps to chromosome 3R, probably to polytene division 32 or 33. Thus the relative contribution of any QTL to oocyst encapsulation varies with the species of parasite. Further, different QTLs were most readily identified in different F2 families. This, like the F1 data, suggests that L3-5 is not genetically homogeneous and that somewhat different pathways may be used to achieve an encapsulation response.

**Conclusion:** We have shown here that different QTLs are involved in responses against different *Plasmodium* parasites.

### Background

Melanotic encapsulation of invading parasites and parasitoids is a widespread insect defense mechanism [1-4]. Depending on the insect and the parasite involved, this

can be either a cellular or a humoral process. The humoral form of melanotic encapsulation occurs primarily, if not exclusively in the order Diptera, and is correlated with low numbers of circulating hemocytes. Melanotic

encapsulation of malaria parasites was first observed within a year of the discovery that mosquitoes are the vectors of malaria [5]. Since then, melanotic encapsulation of plasmodia has been observed in a wide variety of mosquito species. Depending on the mosquito and the parasite species, encapsulation is directed against different stages of development [6].

*Anopheles gambiae* is the principal vector for human malaria in Africa. A laboratory-selected strain derived from the G3 colony of *A. gambiae* was found to deposit melanin capsules around transforming ookinetes or very early oocysts of a variety of *Plasmodium* species [7–10]. Since hemocytes do not appear to participate in formation of these capsules, the mechanism underlying encapsulation is likely to be humoral and/or to involve the midgut epithelium or its basal lamina. One locus, *Pif-B*, was postulated to control encapsulation of *P. cynomolgi* B. Comparisons between mosquitoes from this original refractory strain and a counter-selected susceptible strain showed that refractoriness was highly correlated with the 2L<sup>+</sup> chromosomal arrangement and esterase A. Susceptibility was correlated with inversion 2La and esterase C. A similar correlation was found for the parental G3 colony. Thus it was suggested that *Pif-B* and the esterase A/C polymorphism are located within or very close to the 2La region (divisions 23–26 of ovarian nurse cell polytene chromosome 2L) [11–13]. Genetic mapping experiments with the current refractory (L3-5) and susceptible strains (4Arr) revealed three QTLs controlling the encapsulation of *P. cynomolgi* B. These loci were named *Pen1-3* for *Plasmodium* encapsulation 1, 2 and 3. *Pen1* is the major QTL whereas *Pen2* and *Pen3* are minor QTLs. The refractory alleles at these loci are largely dominant and the contributions of *Pen1* and *Pen2* to the encapsulation response appeared additive [14]. The *Pen1* QTL is also the major QTL for encapsulation of *P. berghei* and for melanin deposition on Sephadex-CM C-25 beads [15]. *Pen1* has been mapped to chromosome 2R, division 8C/D [16], while *Pen2* and *Pen3* locations remain less precisely defined, on distal chromosome 3L in or near division 43, and on proximal chromosome 2R, possibly in division 14, respectively [17,18]. None of these loci affects the intensity of *Plasmodium* infection as measured by number of oocysts on the midgut. Recently, the two loci on chromosome 2L were found to control the *P. falciparum* infection intensity in natural populations of *A. gambiae* [19].

In contrast, refractoriness to *P. cynomolgi* Ceylon in the original refractory strain seemed to be largely but not completely recessive [7,12]. *P. cynomolgi* Ceylon was originally described as a Ceylonese subspecies of *P. cynomolgi* [20], whereas *P. cynomolgi* B originated in Malaysia. The two are currently regarded as being different species (W.E. Collins, personal communication) and molecular data

support this contention [21,22]. Genetic mapping suggested that *Pif-C*, the locus controlling encapsulation of this parasite, assorted independently from the 2La inversion polymorphism, and thus was not linked to the left arm of chromosome 2. However, it was not possible to map this locus further because of the lack of polymorphic markers. More recently, a locus near *Diphenol oxidase-A2* (3R; 33B) on the right arm of the chromosome 3 was also suggested to be important in the melanization of *P. cynomolgi* Ceylon [23]. The refractory allele at this *Dox-A2* linked locus seemed to be most highly expressive when the mosquitoes were heavily infected (geometric mean 178.3, average 212.1 ± SD 104.9 oocysts per midgut). It is not clear whether this locus is the same as *Pif-C*. It is definitely not *Pen2*. Further, it was suggested that a locus or loci unlinked to *Dox-A2*, but also likely to be on chromosome 3, was involved in refractoriness at lower infection intensity, (geometric mean 28.3, average 40.6 ± SD32.7 oocysts per midgut), and may have contributed to parasite encapsulation at high infection intensity as well.


In this study we have investigated the genetics of the encapsulation response against *P. cynomolgi* Ceylon in crosses between the extant refractory L3-5 and susceptible 4Arr strains using microsatellite markers spanning the genome and QTL analysis.

## Results

### Refractoriness to *P. cynomolgi* Ceylon is incompletely recessive

Infection of parental line mosquitoes showed that L3-5 refractory strain mosquitoes encapsulated *P. cynomolgi* Ceylon, whereas 4Arr did not (Fig. 1A). In three separate infection experiments on a total of 54 individual L3-5 mosquitoes, the overall infection prevalence was 81.5%. The number of oocysts in infected individuals ranged between 1 and 147. All oocysts were encapsulated. The infection intensities were generally low and quite variable. The average (± standard deviation) of the intensity of infection varied from a low of 4.93 ± 2.62, to a high of 47.80 ± 40.95 between infections. The overall intensity of infection was 26.36 ± 34.09.

In parallel infection experiments on a total of 60 4Arr susceptible strain mosquitoes blood-fed on the same monkeys used to feed the refractory ones, the infection prevalence was 70.0%. Despite the apparent difference in infection prevalence, the overall intensity of infection was very similar to that of the L3-5 females, 21.50 ± 19.66. Further, the lowest and highest infection intensities, 7.00 ± 7.51 and 32.94 ± 23.12 in 4Arr corresponded to the lowest and highest infection intensities for L3-5 that fed on the same monkeys on the same days. Oocyst numbers ranged between 1 and 79. One encapsulated oocyst was found in each of two 4Arr mosquitoes. These data suggest



**Figure 1**

Melanotic encapsulation of *P. cynomolgi* Ceylon by *A. gambiae*. (A) Responses of individual parental L3-5 refractory strain (squares and circles) and 4Arr susceptible strain females (diamonds and triangles) to *P. cynomolgi* Ceylon infection. Exp. 1 and exp. 2 indicate two separate infection experiments. (B) Responses of F1 female progeny from one cross between L3-5 females and 4Arr males (closed diamonds) and three crosses between 4Arr females and L3-5 males (open squares, circles and triangles).

that if there are differences in *P. cynomolgi* Ceylon infection intensities, they are more likely to be due to the infection parameters of the vertebrate host than to differences intrinsic to the parental mosquito lines.

Reciprocal crosses between L3-5 and 4Arr yielded F1 progeny that were mostly susceptible by the criterion they had no or a low fraction of encapsulated oocysts (Fig. 1B). When L3-5 females were mated with 4Arr males, 33 tested F1 progeny had an average of 27% encapsulated oocysts (infection intensity of  $12.46 \pm 7.31$ ). The reciprocal cross

yielded 101 F1 progeny with an average of 39% encapsulated oocysts (infection intensity of  $18.33 \pm 17.10$ ). Thus refractoriness to *P. cynomolgi* Ceylon in L3-5 mosquitoes appeared to be largely recessive, but less so than observed in the originally selected strain [24]. Further, these data give no evidence for any maternal effects on the encapsulation response.

#### Phenotypic responses of F2 females against *P. cynomolgi* Ceylon

The encapsulation response of 167 females in six F2 intercross families was tested in three separate feeding experiments (Table 2). Family 8-6 females fed on one infected monkey while females in the other five families fed on a different monkey on two successive days of ascending parasitemia. The infection intensity for the combined F2 progeny was  $22.96 \pm 36.14$ . The majority of the F2 progeny were either highly refractory or highly susceptible (Table 2; Figure 2), with 58 (35%) and 47 (28%) individuals exhibiting  $\geq 80\%$  or  $\leq 20\%$  oocysts encapsulated. 62 (37%) of the females had intermediate fractions of encapsulated oocysts. These data suggest that alleles controlling melanotic encapsulation are segregating at more than one locus.

#### Genotyping and genotype frequencies

Genotyping was performed on these 167 female progeny and their six female F1 parents at 52 microsatellite loci spanning the current genetic map of *A. gambiae*. These included 6, 22, and 24 loci on the X, second and third chromosomes, respectively (see Materials and Methods). Six markers were monomorphic in all families and thus uninformative (*H678* on the X; *H682* and *H117* on the second chromosome; *H776*, *H525*, and *H544* on the third chromosome). Marker *H197* on the second chromosome gave uninterpretable data.

Only two alleles were found for each informative marker on the X chromosome. However, among the six F1 female parents of the F2 families, more than two alleles were found at 11 autosomal loci, indicating that one or both of the parental strains was polymorphic at these loci. The *H750* locus, which maps to 65.7 centiMorgan (cM) on the third chromosome, had four alleles. Furthermore, the two parental strains also share some common alleles. This is exhibited by the observation that one of the six F1 females (12-1) was homozygous at 10 autosomal loci. In total, 5, 16 and 14 loci were informative on the X, second and the third chromosomes, respectively. Genotypes of the F2 female progeny were tested for deviations from the expected Mendelian ratios at each locus within each family. Although significant deviations were found for some loci, these loci were not clustered in any chromosomal region (data not shown).

**Table 1: New microsatellite markers and primers developed for this study.**

Marker name	Genetic (Cytogenetic) location	Forward primer (5' – 3')	Repeat type	Reverse primer (5' – 3')	Sequenced allele size (bp)
<b>Chromosome 2</b>					
H682	(2R, 7B)	CGGGCGGTCGTTATTAGGAG	(AC)7	CTGAGTTGACACGAGTTGCC	73
3F11-C1	(2R,9B)	AGATGGTGTCTTCGGTGTC	(GT)17	GTGGAGGAGAAGCCACACAT	156
3B20-B1	(2L, 24B)	GAGACAAAACATGTCAACCG	(CA)17	GTATGTCTGCGCATGAACGAT	122
R007	71.1 (2L, 26D)	GCACCAGTCAAATACGCACT	(CA)3+9	GTCCGTGTTGTTCCAAGATG	184
<b>Chromosome 3</b>					
H249	11.9 (3R, 30A)	GGAAGCAGCTGCCTTTTATG	(TG)15	ACAATGTAACACAGCGCGAG	161
3A13-12	(3R, 30E)	GAACAAAACGTGCCACAGC	(GT)9	TCCCGACTTACGTGAACC	168
3A7-37	(3R, 31B)	CGATTGGAACCAACAAGTGG	(CA)2+9	AAGCCATCTCTTGACGTTCCG	151
Ddc*	(3R, 33B)	CGTTGAGAACC GAAAACAGG	(GT)10	ATCGGGACACTCAGCAACAG	121
3A6-17	(3R, 33C)	TGTGAAGAATGTTTTCGCTGAC	(AC)17	CTCCCCCGTGAATTTGAAC	120
9D17-42	(3R, 33C)	TAAACCTCACCGTCCACATC	(TG)23+5	AAATTGAAACCCGTCCAAAG	150
6B10-1	(3R, 33D)	CATGAGTGCCGCTGTAATG	(TG)2+16	ATCGCTCTCCTCCTAGCC	120
3A23-100	(3R, 34A)	AATGGACCATCGTTAGCACAG	(TG)12	TCCAAAGTTTTCCCTCCAC	136
3L09-C1	(3L, 38C)	GAAACTTCCTGAATTTCCGC	(CA)10+8	ATGGAGGCGACTAAACGTTG	169
				C	
3A2-29	(3L, 41A)	GGCTGCTCTAGGAGGATGTG	(CT)16	AATCGAGGGTGGTGTGAATG	189
3O10-C2	(3L, 41D)	TTAATCACCCACCTTGATCGC	(CA)18	CCTAGAGCTTGAATCGAATC	101
3A14-11	(3L, 43D)	GTAATAGCGAGCGAGCGAAC	(TG)36	GAGCGATCCGAATCGTTAGT	180
3A14-47	(3L, 43D)	GGAGGCACTCTACCACAACC	(GT)10	CCGAACGTCATCGCTAATG	104
3A19-32	(3L, 43D)	CGTGCTGGCATTAAAAAGTG	(CA)15	CTACGCATGCATCTTTCTCC	138
3A5-25	(3L, 45A)	CCTTGGGCGCTATAAAACAC	(TG)10	TATTAGCAGCAGGGGAAACC	131
19A18-54	(3L, 46A)	TAATGCAGTGGTGC GGTTT	(GT)5+6	AATCACAGCCCAGTCAGTCC	121
19C20-B1	(3L, 46B)	ACCAAGTGCTCTTTGGGTTG	(TG)15	ACGAATGCTGGACAATGTGA	141

The cytogenetic location of H682 is *A. gambiae* scaffold predicted by its location in AAAB01008987, contig 566 (GenBank accession AAAB01008987). New primer pairs were designed for markers R007 [44] and H249 [14]. Sequenced allele size is expressed in base pairs (bp). \* This *Dopa decarboxylase* microsatellite is contained within the large first intron of predicted Ddc isoform I. DNA sequence is from G3 strain (GenBank # AF063021; P. Romans, unpublished).


**Table 2: Prevalence and intensity of oocyst infections in the F2 progeny**

Family	Total # of F2 female progeny	Prevalence*	Infection range#	Average ± SD	Parasite/μl †
8-3	24	0.67	1 – 152	35.25 ± 39.73	34,750
8-4	20	0.75	3 – 76	23.8 ± 24.28	34,750
8-6	29	0.55	2 – 30	8.50 ± 7.22	19,080
10-1	38	0.97	3 – 149	44.89 ± 34.42	34,750
12-1	31	0.65	9 – 180	65.15 ± 44.38	34,750
13-4	25	0.76	4 – 104	41.11 ± 26.78	34,750
Total	167	0.74 (123/167)	1 – 180	39.04 ± 36.14	

\* Fraction of F2 females that became infected. # Number of oocysts in the infected midguts. † Number of parasites per microliter of blood at the time of feeding.

**Quantitative trait loci for encapsulation of *P. cynomolgi* Ceylon**  
 No QTL was identified for infection intensity in the previous study [14]. Nor did our infection data suggest any genetic variation for infection intensity between L3-5 and 4Arr. We thus focus our attention on the melanotic encapsulation phenotype, which we have expressed as fraction

of encapsulated oocysts in a single mosquito. Linear regression analysis was performed for each locus within each family and for all families combined. Considering the combined data, no significant association was found between the oocyst encapsulation phenotype and any marker on the X chromosome (Table 3). This is similar to



**Figure 2**

*P. cynomolgi* Ceylon infection phenotypes of F2 female progeny from crosses of six F1 females with their sibling males. (A) Intensity of infection was plotted against the total number of oocysts, both normal and encapsulated, per female midgut. N = 167. (B) Encapsulation responses of 123 infected F2 females. The number of F2 females was plotted against the fraction of oocysts encapsulated per midgut.

previous results obtained for *P. cynomolgi* B and all other parasites tested.

Surprisingly, no major QTL was identified on chromosome 2 near *H290* or *H175* (Table 4), the markers that most precisely defined *Pen1* [14]. Thus it is unlikely that a *Pen1* allele has a significant role in the encapsulation of *P. cynomolgi* Ceylon. *3F11-C1* (2R, 9B) was the only marker near the *Pen1* cytogenetic region that might be associated with an encapsulation QTL. *3F11-C1* had a very high average maximum *F* statistic of 200.88 and gave an *F* statistic calculated from data on all informative individuals of

11.74, though the *P* value associated with it, 0.16, is not significant. More proximally, *H135* and the closely linked marker *H095* appear to define a QTL, which is termed *Pcen2R* (for *P. cynomolgi* Ceylon encapsulation, chromosome arm 2R). These two markers are the same markers most closely associated with *Pen3*, a minor QTL for *P. cynomolgi* B encapsulation. *H095* exhibited the highest average maximum *F* statistic of this data set, 287.99, with an overall maximum of 12.70, the highest of any marker on chromosome 2. However these overall maximum *F* statistics were lower than similar statistics associated with many chromosome 3 loci (Table 5).

There are at least two encapsulation QTLs on chromosome 3 (Table 5). Markers *H555/H158* define a novel QTL in the central region of 3R, in division 32 or 33 that we have named *Pcen3R*. Marker *3A2-29* (3L, 41A) defines another QTL on 3L, *Pcen3L*, which covers the region spanned by divisions 39 – 43. This QTL may explain the relatively high *F* statistics exhibited for markers *H750/H127* in division 39 to *H758* in division 43B. It should be noted that *H758* is the marker most closely associated with *Pen2* and that marker *3A2-29* was not available at the time of the *P. cynomolgi* B encapsulation studies.

When the complete set of progeny was analyzed as a whole, the QTLs on 2R, 3R and 3L explained 13%, 16% and 26% of the refractoriness to *P. cynomolgi* Ceylon, respectively. Similar results were obtained when only all the families except 8-6 (which fed on a different monkey with different parasite level) were analyzed (data not shown).

The refractory and susceptible lines were not highly inbred except for malaria refractoriness or susceptibility and *In(2L)+a* (see Methods). This posed a significant challenge to determining the parental origin of alleles at some loci, in some families. In these cases, the phase information of each allele was predicted using a maximum likelihood method. When a Hidden Markov Model was used to determine the number of QTLs based on the derived phase information, it yielded similar results with two QTLs on the third chromosome and one on the second chromosome. These QTLs were linked with the same marker loci as those identified using single locus linear regression analysis (data not shown, S.W and H. Z, unpublished).

*F* statistics were also calculated for each marker within each individual F2 family. The results for all significant QTLs and markers associated with high *F* statistics in any family are presented in Table 6. Data for *H175*, the marker closest to *Pen1*, are included for comparison. There appeared to be considerable F2 family specificity as to

**Table 3: Probability of a QTL for *P. cynomolgi* Ceylon encapsulation on the X chromosome.**

Locus	Genetic (Cytogenetic) Location	Maximum <i>F</i> statistic*	<i>R</i> <sup>2</sup> †	Average of Family Maximum <i>F</i> statistics ‡	<i>P</i> §	<i>P</i> ' §	Average of Family <i>R</i> <sup>2</sup>
H145	1.9 (4C)	0.64	0.0052	0.31	0.956	1.0	0.0194
H503	11.8 (4B)	2.71	0.0217	0.96	0.612	1.0	0.0365
H036	13.1 (4B)	0.37	0.0032	0.94	0.605	1.0	0.0557
H053	25.0 (3D)	4.16	0.0335	2.67	0.041	0.205	0.1040
H711	40.8 (2A)	5.94	0.0495	1.57	0.273	1.0	0.0677

\* Maximum *F* statistic calculated using all informative data. † Calculated from all informative data. ‡ Average of the maximum *F* statistics for all informative families. § Probability values *P* and *P*' were calculated based on 1000 permutation tests before and after Bonferroni correction respectively.

**Table 4: Probability of a QTL for *P. cynomolgi* Ceylon encapsulation on chromosome 2.**

Locus	Genetic (Cytogenetic) Location	Maximum <i>F</i> statistic*	<i>R</i> <sup>2</sup> †	Average of Family Maximum <i>F</i> statistics ‡	<i>P</i> §	<i>P</i> ' §	Average of Family <i>R</i> <sup>2</sup>
H417	0.0 (2R, 7B)	4.19	0.0398	3.48	0.069	1.0	0.1602
H784b	2.1 (2R, 7A – 10D)	2.23	0.0304	2.55	0.37	1.0	0.1639
<u>H175</u>	3.1 (2R, 8C)	6.93	0.0901	5.76	0.039	0.624	0.2347
H290	4.3 (2R, 8C)	0.32	0.0043	3.21	0.167	1.0	0.1733
H799	18.7 (2R, 9B)	3.30	0.0839	3.30	0.074	1.0	0.0839
3F11-C1	(2R, 9B)	11.74	0.1240	200.88	0.01	0.16	0.4640
H187	24.7 (2R, 11A)	6.56	0.0588	5.15	0.024	0.348	0.1755
H095	42.3 (2R, 13B)	12.70	0.1313	287.99	0.001	0.016	0.4184
<u>H135</u>	47.3 (2R, 14D)	9.48	0.0721	258.33	0.000	0.000	0.3389
H770	50.1 (2R, 15D – 16A)	5.85	0.0542	3.06	0.071	1.0	0.1399
3B20-B1	(2L, 24B)	0.13	0.0011	3.14	0.171	1.0	0.1297
H772	52.9 (2L, 25D – 28D)	1.19	0.0097	2.06	0.219	1.0	0.0901
H603	54.7 (2L, 26D)	0.04	0.0004	3.35	0.132	1.0	0.1381
H819	55.1 (2L, 25D – 28D)	2.61	0.0305	3.28	0.143	1.0	0.1572
H802	57.0 (2L, 26D)	1.52	0.0127	4.69	0.027	0.432	0.1606
R007	71.1 (2L, 26D)	1.26	0.0105	3.67	0.109	1.0	0.1578

Conventions are as for Table 3. Underlined markers H175 and H135 indicate maximum LOD scores for *Pen1* and *Pen3* respectively [14]. Cytogenetic map positions have been used to order loci on 2L where possible. Cytogenetic and genetic map positions are derived from [14,44–46] and unpublished. Cytogenetic locations were further confirmed by comparison to the current scaffolds of the *A. gambiae* genome (GenBank accessions AAAB01000001–AAAB01008987). Where the precise location of a marker is not known, its cytogenetic location is shown as the range covered by the corresponding scaffold.

**Table 5: Probability of a QTL for *P. cynomolgi* Ceylon encapsulation on chromosome 3.**

Locus	Genetic (Cytogenetic) Location	Maximum <i>F</i> statistic*	<i>R</i> <sup>2</sup> †	Average of Family Maximum <i>F</i> statistics ‡	<i>P</i> §	<i>P</i> ' §	Average of Family <i>R</i> <sup>2</sup>
H249	11.9 (3R, 30A)	0.40	0.0451	6.03	0.019	0.285	0.2217
3A13-12	(3R, 30E)	2.27	0.0451	1.96	0.18	1.0	0.1087
H158	46.8 (3R, 32D)	5.43	0.1576	8.73	0.005	0.075	0.2690
H555	37.1 (3R, 33B)	5.97	0.0786	7.03	0.002	0.03	0.2343
3A6-17	(3R, 33C)	32.59	0.2208	9.77	0.024	0.36	0.3208
6B10-1	(3R, 33D)	6.36	0.0696	5.40	0.054	0.81	0.2120
H750	68.0 (3L, 39A)	26.59	0.1789	5.73	0.006	0.09	0.2063
H127	68.2 (3L, 39A)	30.20	0.1984	7.14	0.003	0.045	0.2539
3A2-29	(3L, 41A)	31.13	0.2658	30.4096	0.004	0.06	0.4906
H577	71.9 (3L, 42A)	7.74	0.1770	7.74	0.015	0.225	0.1770
<u>H758</u>	81.0 (3L, 43B)	18.24	0.1401	9.2693	0.006	0.09	0.3094
3A19-32	(3L, 43D)	16.22	0.1209	8.8599	0.01	0.15	0.3033
H753	88.7 (3L, 43D – 46D)	9.24	0.1036	7.5450	0.014	0.21	0.2060
3A14-47	(3L, 43D)	7.86	0.1792	7.8612	0.02	0.30	0.1792

Conventions are as for Table 3. Underlined marker H758 indicates maximum LOD score for *Pen2* [14]

**Table 6: F statistics for markers associated with significant QTL loci within each individual family.**

Family	8-3	8-4	8-6	10-1	12-1	13-4
<b>Locus</b>						
<u>H175</u>	0	9.56	0.0	1.15	0	6.56
3F11-C1	4.16*	757.55	42.00	3.30	1.17	0.69
H095	0.89	1408.89	18.75	7.05*	3.16	0.90
<u>H135</u>	8.27	1522.87	8.09	7.05	3.16	0.57
H158	0.003	5.95	17.45	1.23	11.02	1.27
H555	0.05	3.98*	17.45	11.11	9.35	1.28
3A6-17	8.59	3.98	5.80	26.23	8.02	5.99
H750	8.59	5.95	4.69	14.05	1.09	0.04
H127	8.59	5.95	4.69	14.15	9.35	0.23
3A2-29	8.59*	8.03	4.69*	37.62	69.99	5.99
<u>H758</u>	8.27	2.94	8.40	10.19	23.84	1.98
3A19-32	8.27	2.94	9.39	8.78	20.66	3.14

\*The marker was not informative in this family. The F statistic shown was from the cytogenetically closest neighboring informative locus in the same family. Underlined markers H135 and H758 defined *P. cynomolgi* B encapsulation QTLs *Pen3* and *Pen 2* respectively [14]

which QTLs contributed to refractoriness. Much of the H095 F statistic was due to family 8-4.

*Pcen2R* explained the majority of the encapsulation phenotype in family 8-4, less of the phenotype in family 8-6, very little in family 13-4 and contributed marginally to the phenotype in the other three families. *Pcen3R* appeared to be most important in determining refractoriness in family 8-6 and less important in families 10-1 and 12-1. *Pcen3L* on the other hand, appeared to be the major determinant of refractoriness in families 10-1 and 12-1. It probably played a minor role in the other families.

## Discussion

Melanotic encapsulation of *P. cynomolgi* Ceylon parasites is controlled by incompletely recessive alleles in *A. gambiae* L3-5. Similar results were obtained in previous analysis of the original R and S strains with this parasite [12,23]. Refractoriness of L3-5 to *P. cynomolgi* Ceylon is shown here to be controlled by at least three quantitative trait loci that we have named *Pcen2R*, *3R* and *3L*. Two of the QTLs (*Pcen2R* and *Pcen3L*) are located near *Pen3* and *Pen2* respectively, previously identified as minor QTLs involved in the encapsulation response against *P. cynomolgi* B. These loci may actually be *Pen2* and *Pen3*. A third QTL, *Pcen3R*, is most closely associated with microsatellite locus *H158*. This QTL, therefore, is located in the central region of chromosome 3R, in or near division 32 or 33. This is a novel QTL. A nearby locus, *H555*, was associated with a local peak in LOD score in the *P. cynomolgi* B studies [14]. However, in that analysis, this peak in LOD score did not appear as a QTL separate from *Pen2*. The previously identified major QTL for encapsulation of *P. cynomolgi* B, *Pen1*, contributed little to the encapsulation

of *P. cynomolgi* Ceylon. Thus, it appears that different genetic loci may be involved in responses to infections by different malaria parasites or, perhaps more precisely, may make malaria species-dependent contributions to the genetic variance in encapsulation.

Initial studies of *A. gambiae* refractoriness to *P. cynomolgi* Ceylon used non-homosequential mosquito strains in which a major difference between the refractory and susceptible strains was the absence or presence of inversion *2La*. This inversion encompasses polytene chromosome divisions 23 through 26. It was absent in the refractory strain and fixed in the counter-selected susceptible one. Those studies suggested that melanotic encapsulation of this parasite was controlled by a largely recessive allele at one locus, *Pif-C* [12]. This locus assorted independently from *Pif-B*, a major locus for encapsulation of *P. cynomolgi* B that appeared to be tightly associated the *2L+<sup>a</sup>* chromosomal arrangement [11–13]. Vernick et al. [12] also found that a refractory allele at *Pif-B* acted as a dominant enhancer of expressivity of the refractoriness to *P. cynomolgi* Ceylon. A later study using the same R strain and a backcross strategy ensuring that each offspring received at least one refractory allele of *Pif-B* showed that the refractory phenotype is controlled by at least two, largely recessive loci on chromosome 3 [23]. One of these loci appeared linked to *Dox-A2* (3R; 33B). Additional analysis of those data suggests that this locus maps approximately 32 cM from *Dox-A2*. (P. Romans, unpublished). Therefore this locus is expected to be located extremely distally on chromosome 3R, or else to be tightly linked to the centromere on 3R or 3L. The other locus or loci assorted independently of *Dox-A2* but did not assort independently of the *Dox-A2* linked refractori-

ness locus. It was not mapped further because of lack of genetic markers. This study provided strong evidence for variable penetrance and expressivity of refractory alleles at the *Dox-A2* linked locus. As indicated by Romans et al. [23], this locus was more completely expressive when the geometric mean of the parasite load was 178 oocysts per midgut, rather than when it was 28. Since the infection intensities in the present study were similar (geometric mean 22.96, average  $39.04 \pm \text{SD } 36.14$ ) to this lower number, it is possible that the *Dox-A2* linked locus could have escaped detection even though most of the third chromosome was subjected to QTL analysis. Nevertheless, the 3A2-29 QTL in proximal 3L maps at the appropriate distance from *Dox-A2* to be the *Dox-A2*-linked gene. Further, it is linked to *H758* in division 43, a marker that would assort independently of *Dox-A2*. Therefore, if *H758* is not part of the QTL associated with marker 3A2-29, but is a separate QTL, markers 3A2-29 and *H758* could define the loci inferred in the previous study.

Our observation that different genes may be responsible for encapsulation of different parasite species raises the possibility that different QTL(s) might be required for melanotic encapsulation of human malaria parasites, such *Plasmodium falciparum*. The previous selected refractory strain encapsulated *P. falciparum* isolates from Africa less effectively [7]. Whether refractoriness to *P. falciparum* is dominant or recessive has not been determined for the previous selected or the L3-5 refractory strain of *A. gambiae*.

Our observation that different genes may be responsible for encapsulation of different parasite species is not unique. In *Drosophila melanogaster*, a strain that encapsulates eggs of the parasitic wasp *Leptopilina boulardi* L104, a Congolese strain, is susceptible to parasitism by *L. boulardi* G317 from Tunisia and by *L. heterotoma* [25–28]. Further, genetic mapping experiments have shown that the encapsulation responses to *L. boulardi* and *Asobara tabida* are controlled by single dominant, but different loci. The locus for *L. boulardi* encapsulation, *Rlb*, maps to chromosome 2 at approximately 86.7 cM, in the region 55E3 – E6;F3 on the polytene map [27,29,30]. *Rat*, the locus for *A. tabida* encapsulation, also maps to chromosome 2R, but at 51.3 cM [26,30]. The identities of these genes have not yet been determined. Benassi et al., [26] have suggested that the products of these loci should be involved in parasitoid recognition.

In the G3 parental *A. gambiae* strain from which the previous and current inbred refractory strains were derived, >50% of females encapsulate *P. cynomolgi* B, though only 1 – 3 % of females encapsulate *P. cynomolgi* Ceylon. Recently, field-caught *A. gambiae sensu lato* mosquitoes from Tanzania were tested for their ability to melanize *P.*

*falciparum* and CM Sephadex C-25 beads [31]. While fewer than 1% of these mosquitoes were able to encapsulate *P. falciparum* oocysts, 90% melanized the beads. Even though these results with a natural population of mosquitoes could be confounded by species, age, nutritional and reproductive status [32], they are intriguingly similar to results obtained with G3. It is possible that genetic polymorphism in the encapsulation response may be maintained by selection in natural populations.

The identities of the genes contributing to malaria refractoriness in *A. gambiae* are currently unknown. Both susceptible and refractory strains deposit melanin in egg chorions and in the cuticles of larvae and adults. Thus it is likely that refractoriness loci encode variant molecules that somehow expose parasites as foreign, or directly affect the signaling pathways involved in activation of the phenoloxidase cascade. Encapsulation occurs just as ookinetes penetrate through the basal labyrinth of the midgut, contact the basal lamina and become exposed to the contents of the hemocoel [8–10]. It is also possible that defects in the basal lamina could be responsible for exposing parasites to soluble receptors and components of the phenoloxidase cascade.

It is quite possible that soluble receptors are involved in parasite recognition and initiation of encapsulation. Paskewitz and Riehl [33] showed that if Sephadex CM C-25 beads had been exposed first to hemolymph from susceptible *A. gambiae* 4Arr, they could not subsequently be encapsulated by refractory L3-5 mosquitoes. This result could indicate that the receptors are soluble, although other interpretations are possible.

L3-5 refractory mosquitoes appear somewhat darker than 4Arr susceptible ones at all stages beyond L2. Mortality associated with molting from L3 to L4, from L4 to pupa and from pupa to adult is characterized by extreme melanization in the L3-5 strain (P. Romans and C. Brotherston, unpublished). It is unlikely that this difference is due to phenoloxidase enzymes themselves: Brey et al. [34] showed that the original refractory line of *A. gambiae* and its G3 parent did not differ significantly in phenoloxidase activity. Thus it is also possible that some refractoriness loci encode proteins involved in regulating activation of the phenoloxidase cascade, putting it on a sensitive trigger downstream of any receptor, or that they encode enzymes involved in production of phenoloxidase substrates.

## Methods

### Mosquitoes and parasites

Mosquitoes and parasites were maintained at the Centers for Diseases Control and Prevention, Atlanta GA as described previously [35,36]. The derivation of the *A. gambiae* malaria refractory L3-5 and malaria susceptible

4Arr strains used in these crosses is given by [14,37]. Reciprocal crosses between L3-5 and 4Arr were carried out en masse. F1 offspring of each reciprocal cross were sib-mated and females were then allowed to oviposit individually in order to establish F2 families of mosquitoes. F2 females were then allowed to feed on rhesus monkeys infected with *P. cynomolgi* Ceylon. These infections were carried out under the approved protocol 676-COL-MON-CH. Midguts were dissected from F2 female mosquitoes 8 – 10 days post-blood meal and stained with 2% mercurochrome (Sigma, St Louis, USA). The numbers of normal and encapsulated oocysts on each midgut were determined under a microscope [7].

### Microsatellite Analysis

Genomic DNA extraction and genotyping of *A. gambiae* were carried out as described [14,38]. Most of the microsatellite marker alleles were scored using an ALFexpress DNA sequencer (Amersham Biosciences, Piscataway, NJ) with Cy-5 labeled primers from Sigma Genosys (The Woodlands, TX), Genset (La Jolla, CA), or Integrated DNA Technologies (Coralville, IA). Some markers were scored using <sup>32</sup>P-end-labeled primers as described [39]. The previously described markers we employed were H145, H503, H036, H053, H711 and H678 for the X chromosome; H682, H784b, H417, H290, H175, H197, H187, H799, H095, H135, H770, H772, H603, H819, H802, H117, and R007 for chromosome 2; H776, H249, H525, H555, H158, H750, H127, H577, H758, H753, and H544 for chromosome 3 [38,40]. New primer pairs designed for R007 and H249 are listed in Table 1. Additional new microsatellite markers and primer pairs designed for this study are also described in Table 1. With one exception these markers were identified among *A. gambiae* PEST strain bacterial artificial chromosome clones (BACs) from the Notre Dame-1 library (Collins *et al.*, unpublished), referred to as ND-1 by [41], either from end sequences available at Genoscope <http://www.cns.fr/externe/English/Projets/Resultats/rapport.html> or by screening HindIII or EcoRI subclones of BACs (that have been mapped to the chromosome 3 by in situ hybridization) with <sup>32</sup>P-end labeled d(GT)<sub>15</sub> and d(CT)<sub>15</sub>. The *Ddc* microsatellite was deduced from G3 sequence, GenBank # AF063021. Genotypes of the F1 female parents and their F2 female progeny were determined.

### Statistical Analyses

Both parental strains were polymorphic at many microsatellite loci, except for those on the X chromosome. This led to different numbers of alleles being found at different loci. Interval mapping [42] was not used for linkage mapping because the statistical analysis requires knowledge of the *cis*- or *trans*-relationships between alleles at different loci, and this could not always be determined. Encapsulation phenotypes were tested for statistical association with

genotypes of the marker loci within families and for the combined informative data. A single locus association test was performed by fitting a linear regression between the trait value (fraction of oocysts encapsulated) and the genotype at each locus within a family. The significance of each *F* statistic was established by calculating the proportion of 1000 permutation samples exhibiting an *F* statistic as large as or larger than the *F* statistic determined from the observed data both before and after Bonferroni correction. The contribution of a given QTL to the genetic variance was estimated by *R*<sup>2</sup> [43].

### Authors' Contributions

MQB performed genetic crosses and infection studies. Additional genetic markers were developed by PR, CL and LZ. LZ and CL performed genotyping experiments. Statistical analyses were performed by SW, HZ. Writing was carried out primarily by PR and LZ.

### Acknowledgements

We thank Frank Collins and Anton Cornel for provision of cytogenetically mapped BAC clones from the Notre Dame-1 library. We also thank JoAnn Sullivan for excellent technical support and Helen Rodd for additional statistical analysis. This work was supported by NIH grant AI43053 and an award from the Burroughs Wellcome Fund to L.Z. C.L. was supported in part by a grant (Project 990478) from UNPD/World Bank/WHO Special Programme for Research and Training in Tropical Diseases (TDR). Additional support was provided by an operating grant to P.R. from the Natural Sciences and Engineering Research Council of Canada.

### References

1. Götz P and Vey A: **Humoral encapsulation in insects.** *Hemocytic and Humoral Immunity in Arthropods* Edited by: AP Gupta. New York, Wiley and Sons; 1986:407-429.
2. Götz P and Boman HG: **Insect immunity.** *Comprehensive Insect Physiology, Biochemistry and Pharmacology Volume 3.* Edited by: GA Kerkut and LI Gilbert. Oxford, Pergamon Press; 1985.
3. Götz P: **Mechanisms of encapsulation in dipteran hosts.** *Immune Mechanisms in Invertebrate Vectors* Edited by: AM Lackie. Oxford, Oxford University Press; 1986:pp.1-20.
4. Lackie AM: **Immune mechanisms in insects.** *Parasitol Today* 1988, 4:98-105.
5. Ross R: **On some peculiar pigmented cells found in two mosquitoes fed on malarial blood.** *The British Medical Journal* 1897, 18 December: 1786-1788.
6. Zheng L, Benedict MQ, Romans P, Paskewitz SM, Collins FH and Kafatos FC: **Genetics of Encapsulation Response of Anopheles gambiae against Plasmodium Parasites.** *IX International Congress of Parasitology* Edited by: I Tada, S Kojima and M Tsuji. Monduzzi Editore, Bologna, Italy; 1998:103-109.
7. Collins FH, Sakai RK, Vernick KD, Paskewitz S, Seeley DC, Miller LH, Collins WE, Campbell CC and Gwadz RW: **Genetic selection of a Plasmodium-refractory strain of the malaria vector Anopheles gambiae.** *Science* 1986, 234:607-610.
8. Paskewitz SM, Brown MR, Lea AO and Collins FH: **Ultrastructure of the encapsulation of Plasmodium cynomolgi (B strain) on the midgut of a refractory strain of Anopheles gambiae.** *J. Parasitol.* 1988, 74:432-439.
9. Paskewitz SM, Brown MR, Collins FH and Lea AO: **Ultrastructural localization of phenoloxidase in the midgut of refractory Anopheles gambiae and association of the enzyme with encapsulated Plasmodium cynomolgi.** *J. Parasitol.* 1989, 75:594-600.
10. Paskewitz SM and Christensen BM: **Immune responses of vectors.** *The biology of disease vectors* Edited by: B J Beaty and W C Marquardt. Niwot, Colorado, University Press of Colorado; 1996.

11. Vernick KD and Collins FH: **Association of a Plasmodium-refractory phenotype with an esterase locus in Anopheles gambiae.** *Am. J. Trop. Med. Hyg.* 1989, **40**:593-597.
12. Vernick KD, Collins FH and Gwadz RW: **A general system of resistance to malaria infection in Anopheles gambiae controlled by two main genetic loci.** *Am. J. Trop. Med. Hyg.* 1989, **40**:585-592.
13. Crews-Oyen AE, Kumar VK and Collins FH: **Association of two esterase genes, a chromosomal inversion, and susceptibility to Plasmodium cynomolgi in the African malaria vector Anopheles gambiae.** *Am. J. Trop. Med. Hyg.* 1993, **49**:341-347.
14. Zheng L, Cornel AJ, Wang R, Erfle H, Voss H, Ansong W, Kafatos FC and Collins FH: **Quantitative trait loci for refractoriness of Anopheles gambiae to Plasmodium cynomolgi B.** *Science* 1997, **276**:425-428.
15. Gorman MJ, Severson DW, Cornel AJ, Collins FH and Paskewitz SM: **Mapping a quantitative trait locus involved in melanotic encapsulation of foreign bodies in the malaria vector, Anopheles gambiae.** *Genetics* 1997, **146**:965-971.
16. Thomasova D, Ton LQ, Copley RR, Zdobnov EM, Wang X, Hong YS, Sim C, Bork P, Kafatos FC and Collins FH: **Comparative genomic analysis in the region of a major Plasmodium-refractoriness locus of Anopheles gambiae.** *Proc Natl Acad Sci U S A* 2002, **99**:8179-8184.
17. Collins FH, Zheng L, Paskewitz SM and Kafatos FC: **Progress in map-based cloning of Anopheles gambiae genes responsible for encapsulation of malaria parasites.** *Annals Trop. Med. Parasitol.* 1997, **91**:517-521.
18. Collins FH, Saunders RD, Kafatos FC, Roth C, Ke Z, Wang X, Dymbrowski K, Ton L and Hogan J: **Genetics in the study of mosquito susceptibility to Plasmodium.** *Parassitologia* 1999, **41**:163-168.
19. Niare O, Markianos K, Volz J, Oduol F, Toure A, Bagayoko M, Sangare D, Traore SF, Wang R, Blass C, Dolo G, Bouare M, Kafatos FC, Kruglyak L, Toure YT and Vernick KD: **Genetic loci affecting resistance to human malaria parasites in a West African mosquito vector population.** *Science* 2002, **298**:213-216.
20. Dissanaike AS, Nelson P and Garnham PCC: **Two new malaria parasites, Plasmodium cynomolgi ceylonensis subsp. nov. and Plasmodium fragile sp. nov., from monkeys in Ceylon.** *Ceylon J Med Sci (D)* 1965, **14**.
21. Galinski MR, Arnot DE, Cochran AH, Barnwell JW, Nussenzweig RS and Enea V: **The circumsporozoite gene of the Plasmodium cynomolgi complex.** *Cell* 1987, **48**:311-319.
22. Escalante AA, Barrio E and Ayala FJ: **Evolutionary origin of human and primate malaria: evidence from the circumsporozoite protein gene.** *Mol Biol & Evo* 1995, **12**:616-626.
23. Romans P, IV W. C. Black, Sakai RK and Gwadz RW: **A gene effecting malaria refractoriness is linked to Diphenol oxidase A2 on Chromosome 3 of Anopheles gambiae.** *Am. J. Trop. Med. Hyg.* 1999, **60**:22-29.
24. Vernick KD, Collins FH, Seeley DC, Gwadz RW and Miller LH: **The genetics and expression of an esterase locus in Anopheles gambiae.** *Biochem. Genet.* 1988, **26**:367-379.
25. Rizki TM, Rizki RM and Carton Y: **Leptopilina heterotoma and L. boulardi: strategies to avoid cellular defense responses of Drosophila melanogaster.** *Experimental Parasitology* 1990, **70**:466-475.
26. Benassi V, Frey F and Carton Y: **A new specific gene for wasp cellular immune resistance in Drosophila.** *Heredity* 1998, **80**:347-352.
27. Carton Y and Nappi AJ: **Drosophila Cellular immunity against parasitoids.** *Parasitol. Today* 1997, **13**:218-227.
28. Zheng L: **Genetic basis of encapsulation response in Anopheles gambiae.** *Parassitologia* 1999, **41**:181-184.
29. Hita MT, Poirie M, Leblanc N, Lemeunier F, Lutcher F, Frey F, Periquet G and Carton Y: **Genetic localization of a Drosophila melanogaster resistance gene to a parasitoid wasp and physical mapping of the region.** *Genome Res* 1999, **9**:471-481.
30. Poirie M, Frey F, Hita M, Huguet E, Lemeunier F, Periquet G and Carton Y: **Drosophila resistance genes to parasitoids: chromosomal location and linkage analysis.** *Proc R Soc Lond B Biol Sci* 2000, **267**:1417-1421.
31. Schwartz A and Koella JC: **Melanization of plasmodium falciparum and C-25 sephadex beads by field-caught Anopheles gambiae (Diptera: Culicidae) from southern Tanzania.** *J Med Entomol* 2002, **39**:84-88.
32. Chun J, Riehle M and Paskewitz SM: **Effect of mosquito age and reproductive status on melanization of sephadex beads in Plasmodium-refractory and -susceptible strains of Anopheles gambiae.** *J Invertebr Pathol* 1995, **66**:11-17.
33. Paskewitz SM and Riehle MA: **Response of Plasmodium refractory and susceptible strains of Anopheles gambiae to inoculated Sephadex beads.** *Dev. Comp. Immunol.* 1994, **18**:369-375.
34. Brey PT, Ahmed A, Lee W-J, Ashida M and Lehane MJ: **Tyrosinase-type prophenoloxidase distribution in the alimentary canal of strains of Anopheles gambiae refractory and susceptible to Plasmodium infection.** *Exp. Parasitol.* 1995, **80**:654-664.
35. Benedict MQ: **Care and maintenance of anopheline mosquito colonies.** *The Molecular Biology of Insect Disease Vectors* Edited by: J M Crampton C B Beard C Louis. London, Chapman and Hall; 1997:3-12.
36. Collins WE: **Infection of mosquitoes with primate malaria.** *The Molecular Biology of Insect Disease Vectors* Edited by: J M Crampton C B Beard C Louis. London, Chapman and Hall; 1997:92-100.
37. Gorman MJ and Paskewitz SM: **A genetic study of a melanization response to Sephadex beads in Plasmodium-refractory and -susceptible strains of Anopheles gambiae.** *Am J Trop Med Hyg* 1997, **56**.
38. Wang R, Kafatos FC and Zheng L: **Microsatellite markers and genotyping procedures for Anopheles gambiae, a malaria vector in Africa.** *Parasitol. Today* 1998, **15**:33-37.
39. Zheng L, Collins FH, Kumar V and Kafatos FC: **A detailed genetic map for the X chromosome of the malaria vector, Anopheles gambiae.** *Science* 1993, **261**:605-608.
40. Zheng L, Benedict MQ, Cornel AJ, Collins FH and Kafatos FC: **An integrated genetic map of the African human malaria vector mosquito, Anopheles gambiae.** *Genetics* 1996, **143**:941-952.
41. Hong YS, Hogan JR, Wang X, Sarkar A, Sim C, Loftus BJ, Ren C, Huff ER, Carlisle JL, Black K, Zhang HB, Gardner MJ and Collins FH: **Construction of a BAC library and generation of BAC end sequence-tagged connectors for genome sequencing of the African malaria mosquito Anopheles gambiae.** *Mol Genet Genomics* 2003, **268**:720-728.
42. Lander E, Green P, Abrahamson J, Barlow A, Daley M, Lincoln S and Newburg L: **Mapmaker: an interactive computer package for constructing primary genetic linkage maps of experimental and natural populations.** *Genomics* 1987, **1**:174-181.
43. Rice JA: **Mathematical statistics and data analysis.** 2nd edition. Belmont, CA, Duxbury Press (Wadsworth Publishing Company); 1995.
44. Dimopoulos G, Zheng L, Kumar V, dellaTorre A, Kafatos FC and Louis C: **Integrated genetic map of Anopheles gambiae: use of RAPD polymorphisms for genetic, cytogenetic and STS landmarks.** *Genetics* 1996, **143**:953-960.
45. Zheng L: **Molecular approaches to mosquito-parasite interactions.** *Arch. Insect Biochem. Physiol.* 1996, **34**:1-18.
46. Wang R, Zheng L, Toure YT, Dandekar T and Kafatos FC: **When genetic distance matters: measuring genetic differentiation at microsatellite loci in whole-genome scans of recent and incipient mosquito species.** *Proc Natl Acad Sci U S A* 2001, **98**:10769-10774.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:  
http://www.biomedcentral.com/info/publishing\_adv.asp

