

Research article

Open Access

A microsatellite linkage map of *Drosophila mojavensis*Regina Staten^{1,2}, Sheri Dixon Schully¹ and Mohamed AF Noor*¹

Address: ¹Department of Biological Sciences, Louisiana State University, Baton Rouge, LA 70803 USA and ²Pennington Biomedical Research Center, Louisiana State University, Baton Rouge, LA 70808 USA

Email: Regina Staten - rstaten@lsu.edu; Sheri Dixon Schully - sdixon1@lsu.edu; Mohamed AF Noor* - mnoor@lsu.edu

* Corresponding author

Published: 26 May 2004

Received: 12 March 2004

BMC Genetics 2004, 5:12

Accepted: 26 May 2004

This article is available from: <http://www.biomedcentral.com/1471-2156/5/12>

© 2004 Staten et al; licensee BioMed Central Ltd. This is an Open Access article: verbatim copying and redistribution of this article are permitted in all media for any purpose, provided this notice is preserved along with the article's original URL.

Abstract

Background: *Drosophila mojavensis* has been a model system for genetic studies of ecological adaptation and speciation. However, despite its use for over half a century, no linkage map has been produced for this species or its close relatives.

Results: We have developed and mapped 90 microsatellites in *D. mojavensis*, and we present a detailed recombinational linkage map of 34 of these microsatellites. A slight excess of repetitive sequence was observed on the X-chromosome relative to the autosomes, and the linkage groups have a greater recombinational length than the homologous *D. melanogaster* chromosome arms. We also confirmed the conservation of Muller's elements in 23 sequences between *D. melanogaster* and *D. mojavensis*.

Conclusions: The microsatellite primer sequences and localizations are presented here and made available to the public. This map will facilitate future quantitative trait locus mapping studies of phenotypes involved in adaptation or reproductive isolation using this species.

Background

Evolutionary biologists have struggled to determine the number and types of genetic changes that lead to speciation. Recent advances in molecular techniques facilitate a more thorough investigation into these issues. For example, by mapping quantitative trait loci (QTLs) affecting interesting traits, we can explore the genetic basis of phenotypic variation between two populations that may lead to reproductive isolation.

One hallmark species used in studies of speciation and ecological adaptation is the desert cactophilic *Drosophila mojavensis*. *D. mojavensis* belongs to the *mulleri* complex of the *repleta* species group within the subgenus *Drosophila*. Unlike many well-studied *Drosophila*, its ecological niche has been well documented, and extensive cytogenetic work has been done on it and its close relative,

D. arizonae [see e.g., [1]]. With regard to speciation, *D. mojavensis* has been the subject of many genetic and phenotypic studies of mate choice [e.g., [2-5]], hybrid sterility and inviability [6,7], and variation in sperm and female sperm-storage organ length [8,9]. However, all of these studies have been forced to use a handful of either allozyme or morphological mutant markers. Microsatellites have been isolated from this species before [10], but they are unmapped and their sequences are not available.

Here, we present a microsatellite-based linkage map of the five major chromosomes of *D. mojavensis* using a new set of markers. We mapped 25 microsatellites to the X chromosome and 65 microsatellites spanning the four major autosomes. We also use our results to confirm the conservation of Muller's chromosome elements [11] across approximately 65 million years of evolutionary

divergence between *D. melanogaster* and *D. mojavensis* [see [12]]. Muller [11] had suggested that chromosomal elements conserve their identities (ie, complement of genes) across all *Drosophila* species, and several subsequent studies have supported this idea [e.g., [13-15]], though only one study involving the *repleta* group [16].

Results and Discussion

Primers were successfully developed for a total of 116 markers. Of these, 26 did not distinguish between the two isofemale lines that were used for mapping and were therefore not pursued. We mapped 25 microsatellites onto the X-chromosome, 10 onto chromosome 2, 7 onto chromosome 3, 13 onto chromosome 4, and 10 onto chromosome 5. Twenty-five more microsatellites were confirmed to be autosomal but could not be mapped because of segregating polymorphism within our lines. Microsatellites were named based on their localizations, where the fifth character of the name was an X if X-linked, A if unmapped autosomal, or a number indicating a specific autosome. The distribution of microsatellites across the chromosomes suggests a possible excess of repetitive sequences on the X-chromosome (27.8% observed vs. 20% expected assuming all chromosomes are similar in size, chi-square test, $p = 0.07$; 27.8% observed vs. 18.7% expected assuming chromosomes are the same size as *D. melanogaster* homologous chromosome arms, chi-square test, $p = 0.03$).

Using two female-parent backcrosses, we constructed a recombinational map of the *Drosophila mojavensis* genome using 34 of our microsatellites: 13 on the X-chromosome, 7 on chromosome 2, 4 on chromosome 3, 7 on chromosome 4, and 3 on chromosome 5. Recombinational distances are presented in Figure 1. *DMOJX040* was not placed in the figure because it was only 0.7 cM from *DMOJX030*. The recombinational lengths of the chromosomes generally exceed the homologous chromosome arms in *D. melanogaster* and some other *Drosophila* species. For example, the X-chromosome in *D. mojavensis* spanned 130.8 cM, while the X-chromosome in *D. melanogaster* spans only 73 cM. Even within the *repleta* species group, *D. buzzatii* has an X-chromosome that spans 109 cM [17] and *D. hydei*'s X spans 116 cM [12]. Similarly, *D. mojavensis* chromosome 2 could only be assembled into three pieces that recombine freely from each other. This difference between species in recombinational length most likely indicates an overall greater recombination rate per megabase in *D. mojavensis*, but we cannot exclude dramatic differences in sequence lengths of the chromosome arms.

We recombinationally mapped some markers in a second cross because of segregating variation within the lines. Figure 1 presents the most conservative map, where all mark-

ers were mapped against each other for any particular chromosome. However, we have some additional information about the linkage of other microsatellites. Specifically, we observed that *DMOJ4200* is freely recombining from all the 4-chromosome markers between and including *DMOJ4010* and *DMOJ4060*. Also, the following markers are freely recombining from each other on chromosome 5: *DMOJ5100*, *DMOJ5200*, *DMOJ5300*, and *DMOJ5400*.

To evaluate the conservation of Muller's elements across 65 million years, we used BLAST [18] to identify segments homologous to the sequences flanking the 65 microsatellites in *D. mojavensis* that were mapped to chromosome. We identified segments mapped to *D. melanogaster* chromosomes similar to 23 of the sequences isolated from *D. mojavensis* (see Table 1). The inferred homology of the arms are as follows (*melanogaster*:*mojavensis*): X:X, 2L:3, 2R:5, 3L:4, 3R:2 [19,20]. Based on the BLAST results, all 23 *D. mojavensis* sequences matched *D. melanogaster* sequences on the homologous chromosome arms. This observation strongly supports the conservation of Muller's elements between the subgenera *Drosophila* (*D. mojavensis*) and *Sophophora* (*D. melanogaster*).

Conclusions


We have developed and mapped a panel of 90 variable microsatellites for genetic studies in a model system for ecological genetics and speciation: *Drosophila mojavensis*. Thirty-four of these microsatellites have been placed onto a detailed linkage map of this species. We also confirmed that Muller's chromosome elements were conserved between *D. melanogaster* and *D. mojavensis*, species separated by 65 million years of independent evolution, in 23 of 65 sequences tested. Given the long-term interest in this species for studies of adaptation and speciation, the construction of a linkage map and presentation of variable microsatellite sequences will facilitate future work in this area.

Methods

Isolation of microsatellite sequences

We used a modification of Hamilton et al's [21] enrichment technique to increase the proportion of microsatellites in the genomic DNA insert library prior to cloning [see also [22]]. This procedure uses a subtractive hybridization, in which streptavidin-coated magnetic beads and biotinylated oligonucleotide repeats retain single-stranded genomic DNA fragments containing repeat sequences.

Genomic DNA was isolated from approximately 30 *D. mojavensis* individuals from a mixture of strains using the Puregene™ DNA Isolation Kit (Genta Systems). Except where indicated, we used reagent concentrations and

**Figure 1**

Linkage map of the five major chromosomes of *Drosophila mojavensis*. From left to right, are the X-chromosome, chromosome 2, chromosome 3, chromosome 4, and chromosome 5. Kosambi recombinational distances between markers are on the left of each chromosome, and the microsatellite names are on the right. A question mark appears between markers or groups when markers were assigned to the same chromosomes but freely recombined from each other.

Table I: Ninety microsatellites mapped in *Drosophila mojavensis*. Microsatellites assigned to chromosome "A" were autosomal but could not be mapped to a particular autosome because of variation segregating within the lines used for mapping. We present the BLAST expect (E) value in the column after the microsatellite name only for the 23 microsatellites used in the Muller's chromosome element comparison.

Name	BLAST E-value	Chromosome	Size	Primers	Repeat motif	GenBank Accession
DMOJX010		X	132	attgtgtcgcccttagggc tgataatttgtgatttgggtgcac	(ca) ₁₁	AY578823
DMOJX020		X	142	ctctgcccactaacggacc ctcttcagtgtgcctttaggtatac	(ca) ₇	AY578824
DMOJX030		X	124	aagctatgccttagtgtcactcc caaacggcattcatataagaatctatctcac	(ag) ₉	AY578825
DMOJX040		X	147	aggcatgccttagttgtgcac cacacatattaagcattgtattacaatcggtcc	(ac) ₁₅	AY578826
DMOJX050		X	143	accaagccaaaagccaattgcac caaagcttgcggcatttag	(ac) ₁₂	AY578827
DMOJX060		X	128	caatttgtggatgtgcgttgcac cagcatttcgtattgcattacac	(ac) ₁₂	AY578828
DMOJX070	1e-19	X	163	gccactcggtgtgcctta atagtctttgtcttatgtcggtg	(ca) ₄	AY578829
DMOJX080		X	202	ccactccacgttcgcgttgc cacaatgttagagcgtcccttaatctcc	(ac) ₁₄	AY578830
DMOJX090		X	149	ccttcaaccttgcgtccagac caatcaaagcgaacgttagctcatg	(ca) ₉	AY578831
DMOJX100		X	160	tttgtatataaggcggaaaggcgg gtctgtatataattataatgttacgttacaaaagaactcac	(gt) ₂₀	AY578832
DMOJX110	2e-09	X	98	cagtggcgcctaaaggctgt gggatgtatgggtctatgggtg	(ca) ₉	AY578833
DMOJX120		X	110	gcctgcatttgtgcattctgc caagtgtttgc当地agctcgag	(tg) ₁₃	AY578834
DMOJX130		X	130	tggctacacttcagcaaaacattc tcatgtgc当地aggttagccagc	(ca) ₁₁	AY578835
DMOJX500		X	100	caattattgtatggccacccc cgagcactttcaatttttgc	(gt) ₁₁	AY578836
DMOJX501		X	136	ctcgagc当地attttattgttgc ccaacgagccatcccttc当地	(ca) ₆	AY578837
DMOJX502		X	107	attaaatgtatccatggacacagccac ctctgc当地tgc当地	(ca) ₁₀	AY578838
DMOJX503		X	136	gccaaggatgtcaatacccttgc aaggcaatatacacaacatcatgtgc	(gt) ₁₀	AY578839
DMOJX504		X	126	catcaatctctagaatgc当地c gagtgtactttaaggc当地gtc	(gt) ₈	AY578840
DMOJX505	0.004	X	128	tcgcttgc当地tgc当地tgc tacgggtatgtc当地atgc	(ac) ₁₀	AY578841
DMOJX506	2e-15	X	149	actgc当地tactgtct acaggctt当地acatggccaaataac	(tc) ₁₆	AY578842
DMOJX507		X	166	ttttctttgc当地tgc当地tgc cacatgtgg当地atgc当地	(tc) ₁₅	AY578843
DMOJX508	4e-43	X	248	aaggc当地tgc当地tgc当地 cattgg当地atgtatgc当地	(ac) ₁₂	AY578844
DMOJX509		X	157	agctgat当地tgc当地tgc当地 ttccat当地tgc当地tgc当地	(ga) ₁₈	AY578845
DMOJX510		X	169	atttggctgtcgccg gattttgtccactgtgc当地	(ag) ₁₈	AY578846
DMOJX511		X	177	gcttc当地tgc当地tgc当地 tgc当地tgc当地tgc当地	(ct) ₁₆	AY578847
DMOJ2010	2		157	acgagtttgc当地tgc当地tgc aaaggcc当地acttgc当地tgc当地	(ac) ₁₄	AY578848
DMOJ2020	2		153	attgactt当地tgc当地tgc当地 cgctgtctc当地tgc当地tgc当地	(gt) ₇	AY578849
DMOJ2030	2		199	tgacgc当地tgc当地tgc当地 gattcaagg当地tgc当地tgc当地	(gt) ₁₆	AY578850
DMOJ2100	4e-05		187	ggcgctcccttaatcacagatac agcatgtgtctgttgc当地	(ag) ₁₇	AY578851

Table I: Ninety microsatellites mapped in *Drosophila mojavensis*. Microsatellites assigned to chromosome "A" were autosomal but could not be mapped to a particular autosome because of variation segregating within the lines used for mapping. We present the BLAST expect (E) value in the column after the microsatellite name only for the 23 microsatellites used in the Muller's chromosome element comparison. (Continued)

DMOJ2200	3e-21	2	148	gtcgctccatagacttacaaggtt cgccccaagaattcacgaagc	(gt) ₉	AY578852
DMOJ2210		2	139	cccagaatgtactctactaag tgctcatcaataaagaaggaaac	(ca) ₈	AY578853
DMOJ2220	0.002	2	136	gttggcttggattggactgag tgtcaatgtgactggcaactg	(gt) ₆	AY578854
DMOJ2300	1e-05	2	114	aattgacagcactccgtggc gttcagcggccgcctac	(ca) ₁₂	AY578855
DMOJ2301		2	158	ctcttagccggagggtcaag aatcttatcgaaaatatgcacacgatgg	(caa) ₁₁	AY578856
DMOJ2302	8e-17	2	193	ctctcgctttctctgtcttatac aactgattaccgctgcatacag	(ac) ₉	AY578857
DMOJ3010	4e-05	3	218	gcccgccggaggtaat atgtgtatggccagtcatattt	(ct) ₈	AY578858
DMOJ3020	4e-23	3	94	acgtggatatacgacgagc tttggccaaatttgagcaactgc	(ca) ₁₄	AY578859
DMOJ3030		3	91	cctagtttctttggccaccc tac cgcagtgaaacgcataggaaaac	(ca) ₁₁	AY578860
DMOJ3040		3	94	gtcagggtcagcagcagc gcctcaacagcacctactgag	(ac) ₁₂	AY578861
DMOJ3100		3	87	ctgatttgtcaccacaggactc gctaattcaaggcacacatgtttag	(ca) ₁₀	AY578862
DMOJ3101	7e-25	3	150	aacggcggcatccgttgc actgtcatcgacaaatgatttta	(gt) ₁₂	AY578863
DMOJ3102		3	210	ctctctgttagaaaaaggctttttaacc tgctgtgtcagcacaaac	(ca) ₁₁	AY578864
DMOJ4010		4	90	agccagtgcaatgcaggc gcctggaccccttgggg	(ca) ₁₆	AY578865
DMOJ4020		4	121	cagcagctgccttattgc aataaatcgccaggcaggac	(ac) ₁₀	AY578866
DMOJ4030	7e-05	4	137	gtatgtttgttaggcacgcata aatgagaatgagaactggacagg	(ca) ₁₀	AY578867
DMOJ4040	1e-07	4	161	gcaacatgtgcctcaactgc ttttccccacacttcttgcagcag	(ca) ₁₁	AY578868
DMOJ4050		4	196	atcgcatagaagacactcat ctggggcaaggaaatgttgc	(ca) ₉	AY578869
DMOJ4060		4	211	cgagactcgctgataagta gattgtattttggccgtgc	(ac) ₄₁	AY578870
DMOJ4100		4	126	cgcagacatatttgtctcc ttcgtagccaaagacaaactcaca	(ga) ₁₁	AY578871
DMOJ4200	3e-10	4	120	gtttcaaggccctgtgatttt caagaagaacaacgcgttatgc	(ct) ₂₄	AY578872
DMOJ4300		4	165	ggaaagaataccaaacgc gtccgcagacagccgc	(ca) ₁₂	AY578873
DMOJ4301		4	133	acatttgctgttaccc ccaatgcctgttgcattatgc	(ca) ₁₂	AY578874
DMOJ4302	2e-42	4	218	gttgtgtcggtggatgttt gacagcactgaaacgatata gatgcata	(ag) ₁₈	AY578875
DMOJ4303		4	174	cacggcaacacttgc ccattgctcatgcgcgttacc	(ag) ₂₃	AY578876
DMOJ4304		4	171	ggcacattgccaca tctgtgcggaaatgc	(ga) ₂₀	AY578877
DMOJ5010	6e-04	5	117	ggcataggacgc gtaaatattcgaaac actacatgc	(ac) ₈	AY578878
DMOJ5020	4e-06	5	144	ctacaggtat acaacagcc	(gt) ₁₁	AY578879
DMOJ5100	8e-18	5	156	agacaactt tgacactgttgc	(gt) ₈	AY578880
DMOJ5200	2e-26	5	154	tgc attttacag at	(ca) ₁₃	AY578881
DMOJ5300		5	97	gttgtggac ttagccaa actttt ttagcata atttagcc	(ca) ₉	AY578882

Table I: Ninety microsatellites mapped in *Drosophila mojavensis*. Microsatellites assigned to chromosome "A" were autosomal but could not be mapped to a particular autosome because of variation segregating within the lines used for mapping. We present the BLAST expect (E) value in the column after the microsatellite name only for the 23 microsatellites used in the Muller's chromosome element comparison. (Continued)

DMOJ5400	5e-08	5	109	cttggatttcagtcagtgcgtc cgccacaatcagtcataagggtcc	(gt) ₁₀	AY578883
DMOJ5500		5	121	ggaaggcgtcactgcatacc gttgtgaaacgtatgttttgcc	(ca) ₁₀	AY578884
DMOJ5501	1e-04	5	99	cgtgccacgtaaacttgc gaaggccaattcaattgtttaggttatccc	(ac) ₉	AY578885
DMOJ5502		5	115	gcataattgacaaggacgtgc tctgagtgcgtcattactttgtatc	(gt) ₁₂	AY578886
DMOJ5503		5	150	gtatacgacatgtggcactgc ttgcaagctggcgtaagc	(tg) ₁₀	AY578887
DMOJA500		A	185	gagacttggacggccgc tcgatagacatgagttttgctagaaacc	(tg) ₈	AY578888
DMOJA501		A	140	tcagtagcctctgcgtacggc cgaacggaaattatgactagtcagcc	(tc) ₃₀	AY578889
DMOJA502		A	138	ctgaaagtctggcagcaagagt gtgttaattttgttttagacgcgtttagag	(ct) ₁₄	AY578890
DMOJA503		A	153	taaggctctgtttcgtaactttgc ctgtcaatgtgttaaacatgtcaacc	(ca) ₉	AY578891
DMOJA504		A	222	aatcatctgcccccccttcac ggaaaatgtgcgtcaggcagggt	(ac) ₁₃	AY578892
DMOJA505		A	181	ccatagtgcgtatgcacgttc gccatagccccatagtagccaaag	(tg) ₁₀	AY578893
DMOJA506		A	147	attaatgcaggccggaaatgtcg gctcgctctgcgtcgatgttatc	(gt) ₁₁	AY578894
DMOJA507		A	134	tcagccggatgttaacttaacttg atgttaccagagcgtatggc	(ac) ₁₂	AY578895
DMOJA508		A	196	ctctgcgcacatgttagactacgc gataaagttaacttttactaccgtatcattc	(tg) ₁₀	AY578896
DMOJA509		A	186	gctgagaacaaaatttcgcgtcc tgttgtgtcccttaacgaaacgttcc	(tg) ₁₈	AY578897
DMOJA510		A	105	cacacagccagacttgcgttag gtttttgattttgtatgcgcattgtcaaac	(tg) ₁₂	AY578898
DMOJA511		A	162	cttttctggcttattacgcggcgc aaaacataatgttaatttgcgtacaagcaac	(tg) ₃₀	AY578899
DMOJA512		A	120	gtatggaaataggcgttgcgtcc gcatatgtatgcggcgttgcgtgc	(ca) ₁₁	AY578900
DMOJA513		A	120	gctcagctaacagaaacaccca gccgtatgcgtcgcgtatct	(tg) ₁₃	AY578901
DMOJA514		A	125	atggcgcacttcgtc gcgcacatgtgcgtcg	(gt) ₁₂	AY578902
DMOJA515		A	203	gaccgaacagcgcgc cacaacactacataaacaccgcgc	(ac) ₁₁	AY578903
DMOJA516		A	163	ggctgttaccaagcacacactc cgctcgatgtgcgtcgatctc	(ca) ₁₃	AY578904
DMOJA517		A	87	gaaaacagctgcaacccgtaaag gctctttaagcgctcaactatatac	(ca) ₁₅	AY578905
DMOJA518		A	118	gtatgtatggcatacagcgg cttggttttatgtatgtatgcgtgtct	(ca) ₇	AY578906
DMOJA519		A	182	atgaataggaatccgcgc agcgccttgcgtgcctac	(ca) ₁₄	AY578907
DMOJA520		A	164	tttcggcgcagggtgc ttatcgttttacccgcatacatgc	(gt) ₈	AY578908
DMOJA521		A	96	ttttgttttagttttgcgcctaacc ttttccataatttgcgtgtgc	(ac) ₉	AY578909
DMOJA522		A	122	cctttcgactgcctccacaac gtcccactacatattgcgtacagctg	(ca) ₁₁	AY578910
DMOJA523		A	147	gcgttagcacagttggactctc tgtctgcggatgtttagtgcgtaa	(ct) ₂₃	AY578911
DMOJA524		A	135	tcgagagatgcgtc cctgttgcattatgtgggtgc	(ac) ₁₂	AY578912

reaction conditions suggested by Hamilton et al [21]. The enrichment procedure was repeated seven times. For each enrichment, one of the following enzymes was used with *NheI* to digest *Drosophila mojavensis* genomic DNA: *Sau3AI*, *BfuCI*, *RsaI*, *AluI*, or *HpyCH4III*. Linker sequences were ligated to the digested DNA to provide a PCR priming site. We then hybridized the digested, linker-ligated DNA to a biotinylated oligonucleotide repeat motif, either (CA)₁₅ or (AG)₁₅, and recovered the microsatellite-enriched DNA. The DNA was amplified via PCR, and fragments between 300 and 800 bp were recovered from an agarose gel for cloning. We then used the Invitrogen TOPO-TA cloning kit to clone the DNA into plasmids and transform into *E. coli*. We omitted the chemiluminescent screen and used pUC19 primers to amplify *D. mojavensis* DNA inserts directly from colonies. Each 50 μl reaction volume contained 50 mM Tris-HCl (pH 8.3), 20 mM KCl, 1.5 mM MgCl₂, 0.2 mM each dNTP, 0.5 μM pUC forward and reverse primers, and 1.0 unit *Taq* polymerase (AmpliTaq, Perkin Elmer). DNA was added by touching a sterile toothpick to a colony and swirling the toothpick into the reaction mix. We used the following thermal profile: 95°C for 5 min; 30 cycles of 94°C for 60 s; 55°C for 30 s, 55°C for 30 s, 72°C for 30 s; rapid thermal ramp to 40°C. PCR products were sequenced with an ABI Prism® Big Dye™ Terminator Cycle Sequencing Ready Reaction Kit, and products were visualized on ABI sequencers in the LSU Museum of Natural Science, Pennington Biomedical Research Center, or the Department of Biological Sciences' genomics facility.

Fly stocks

Several inbred lines of *D. mojavensis* were tested to determine the most suitable lines for constructing a microsatellite map based on microsatellite allelic differences between strains, bearing the same chromosomal arrangements, and lack of segregating microsatellite alleles. In the end, we selected the lines A993 (Rancho El Diamante, Sonora) and A924 (St. Rosa Mtns., AZ), obtained from Dr. William J. Etges. These lines were further brother-sister mated for 9–12 generations to ensure thorough inbreeding and a reduction of segregating alleles.

Microsatellite assay conditions

We designed two primers for each microsatellite-bearing sequence, one bearing an M13(-29) tail. A 10 μL PCR reaction was then performed using 0.5 μM of each primer, 1.0 μL of dNTPs, 1.0 μL of 10X PCR buffer (100 mM Tris pH 8.3, 500 mM KCl, 15 mM MgCl₂), 0.4 μL of IRDye (LiCor), 1U *Taq* DNA polymerase, and 0.5 μL from a single fly DNA preparation (Puregene). We sometimes added 1.0 μL of 10 mM MgCl₂ to the reaction or more polymerase to optimize the results of the PCR. A touchdown PCR cycle was performed [23], and amplifications were visualized on acrylamide gels on our LiCor DNA analyzer.

Assignment to linkage groups

Virgin females and males of the A993 and A924 lines were crossed and offspring reared. DNA was isolated from the parents and progeny using the Puregene™ DNA Isolation Kit (Gentra Systems). We determined if markers differentiating the lines were X-linked or autosomal by comparing the F₁ males to the F₁ females and parent strains. For X-linked markers, males consistently bore one allele while females consistently bore two. Autosomal markers were further tested using 20 progeny of a male-parent backcross. Because there is no recombination in *Drosophila* males, the offspring all inherited a nonrecombinant chromosome from one of the original lines. By comparing genotypes across the male-parent backcross progeny, markers were assigned to linkage groups. We also used the NCBI Basic Local Alignment Search Tool [BLAST: [18]] to identify putatively homologous sequences in *D. melanogaster*. Sequences bearing an expect (E) value below 0.01 were scored, as E-values are nearly identical to probability (p) values in that range.

Recombinational mapping within linkage groups

Virgin F₁ females (progeny of the cross described in "Assigning to linkage groups") were backcrossed to males of one of the pure lines (A924). To ascertain the recombinational distances between the markers on each chromosome, we genotyped the parents and 200 progeny with each marker previously assigned to a linkage group. Recombinational distances were estimated in Kosambi centiMorgans using Mapmaker [24].

Authors' contributions

RS maintained all fly cultures and performed all reactions and analyses involved in the recombinational mapping of microsatellites. SDS and MAFN produced the microsatellite genomic libraries, sequenced the clones, and designed the primers. All authors contributed to the preparation of this manuscript.

Acknowledgements

This research was supported by National Science Foundation grants 9980797, 0211007, and 0314552, and Louisiana Board of Regents Governor's Biotechnology Initiative grant 005 to MAFN and a Sigma Xi grant-in-aid of research to SDS. We thank William J. Etges for providing fly stocks and moral support, Daniel Ortiz-Barrientos, Christy Henzler, and three referees for constructive comments on this manuscript, and Lisa Burk for technical assistance.

References

1. Ruiz A, Heed WB, Wasserman M: **Evolution of the mojavensis cluster of cactophilic *Drosophila* with descriptions of two new species.** *J Hered* 1990, **81**:30-42.
2. Etges WJ: **Premating isolation is determined by larval substrates in cactophilic *Drosophila mojavensis*.** *Evolution* 1992, **46**:1945-1950.
3. Zouros E: **The chromosomal basis of sexual isolation in two sibling species of *Drosophila*: *D. arizonensis* and *D. mojavensis*.** *Genetics* 1981, **97**:703-718.

4. Wasserman M, Koepfer HR: **Character displacement for sexual isolation between *Drosophila mojavensis* and *Drosophila arizonensis*.** *Evolution* 1977, **31**:812-823.
5. Etges WJ, Ahrens MA: **Premating isolation is determined by larval-rearing substrates in cactophilic *Drosophila mojavensis*. V. Deep geographic variation in epicuticular hydrocarbons among isolated populations.** *Am Nat* 2001, **158**:585-598.
6. Zouros E: **The chromosomal basis of viability in interspecific hybrids between *Drosophila arizonensis* and *Drosophila mojavensis*.** *Can J Genet Cytol* 1981, **23**:65-72.
7. Pantazidis AC, Galanopoulos VK, Zouros E: **An autosomal factor from *Drosophila arizoneae* restores spermatogenesis in *Drosophila mojavensis* males carrying the *D. arizoneae* Y chromosome.** *Genetics* 1993, **134**:309-318.
8. Miller GT, Starmer WT, Pitnick S: **Quantitative genetic analysis of among-population variation in sperm and female sperm-storage organ length in *Drosophila mojavensis*.** *Genet Res Camb* 2003, **81**:213-220.
9. Pitnick S, Miller GT, Schneider K, Markow TA: **Ejaculate-female coevolution in *Drosophila mojavensis*.** *Proc Roy Soc Lond B* 2003, **270**:1507-1512.
10. Ross CL, Dyer KA, Erez T, Miller SJ, Jaenike J, Markow TA: **Rapid divergence of microsatellite abundance among species of *Drosophila*.** *Mol Biol Evol* 2003, **20**:1143-1157.
11. Muller HJ: **Bearings of the *Drosophila* work on systematics.** In *New Systematics* Edited by: Huxley J. Oxford: Clarendon Press; 1940:185-268.
12. Powell JR: **Progress and Prospects in Evolutionary Biology: The *Drosophila* Model.** New York: Oxford University Press 1997.
13. Whiting JH, Piley MD, Farmer JL, Jeffery DE: **In situ hybridization analysis of chromosomal homologies in *Drosophila melanogaster* and *Drosophila virilis*.** *Genetics* 1989, **122**:99-109.
14. Papaceit M, Juan E: **Chromosomal homologies between *Drosophila lebanonensis* and *D. melanogaster* determined by in situ hybridization.** *Chromosoma* 1993, **102**:361-368.
15. Gallego P, Juan E, Papaceit M: **Chromosomal homologies between *Drosophila melanogaster* and *D. funebris* determined by in-situ hybridization.** *Chromosome Res* 1999, **7**:331-339.
16. Ranz JM, Segarra C, Ruiz A: **Chromosomal homology and molecular organization of Muller's elements D and E in the *Drosophila repleta* species group.** *Genetics* 1997, **145**:281-295.
17. Schafer DJ, Fredline DK, Knibb WR, Green MM, Barker JSF: **Genetics and linkage mapping of *Drosophila buzzatii*.** *J Hered* 1993, **84**:188-194.
18. Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ: **Basic local alignment search tool.** *J Mol Biol* 1990, **215**:403-410.
19. Wasserman M: **Evolution of the repleta group.** In *The Genetics and Biology of *Drosophila**, 3b Edited by: Ashburner M, Carson HL, Thompson JN. London: Academic Press; 1982:61-139.
20. Wasserman M: **Cytological evolution of the *Drosophila repleta* species group.** In *Drosophila Inversion Polymorphism* Edited by: Krimbas CB, Powell JR. Boca Raton, FL: CRC Press; 1992:455-552.
21. Hamilton MB, Pincus EL, DiFiore A, Fleischer RC: **Universal linker and ligation procedures for construction of genomic DNA libraries enriched for microsatellites.** *BioTechniques* 1999, **27**:500-507.
22. Kandpal RP, Kandpal G, Weissman SM: **Construction of libraries enriched for sequence repeats and jumping clones, and hybridization selection for region-specific markers.** *Proc Natl Acad Sci USA* 1994, **91**:88-92.
23. Palumbi SR: **Nucleic Acids II: The Polymerase Chain Reaction.** In *Molecular Systematics* Edited by: Hillis DM, Moritz C, Mable BK. Sunderland, Mass.: Sinauer Associates, Inc; 1996:205-247.
24. Lander ES, Green P, Abrahamson J, Barlow A, Daly MJ, Lincoln SE, Newburg L: **MAPMAKER: An interactive computer package for constructing primary genetic linkage maps of experimental and natural populations.** *Genomics* 1987, **1**:174-181.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

